

## COMMENT ON “MEANING OF THE WAVE FUNCTION”

J.Samuel and R.Nityananda  
Raman Research Institute  
Bangalore, INDIA 560 080

*Abstract* : We draw attention to an elementary flaw in a recently proposed experiment to measure the wave function of a single quantum system.

In a recent paper, Aharonov, Anandan and Vaidman [1] claim to have devised an experiment to measure the wave function [2] of a *single* quantum mechanical system. The experiment they propose is a ‘protective measurement’: while the measurement is being performed, the wave function is prevented from collapsing by means of suitably chosen external field. The claim of ref.[1] is at variance with accepted interpretations of quantum mechanics. According to these, it is impossible to learn anything new about a quantum system without disturbing it. Our purpose in this note is to examine the claim made in ref.[1] more closely.

The issues involved are best illustrated by the following simple example. One can determine the state of polarisation of a classical laser beam (one with a large number of photons) by means of simple experiments involving polaroids. Given a single photon, conventional wisdom holds that there is no way to measure its polarisation precisely. If the photon passes through a

polaroid, all one can say with any certainty is that it was not polarised in the direction orthogonal to the polaroid. Ref.[1] challenges this conventional wisdom.

Much of reference [1] addresses the question of whether the wave function has an “epistemological” or “ontological” meaning. It is not our intention to enter into this debate. We restrict our attention to the simplest of the experiments proposed in [1], which is a modified Stern-Gerlach experiment. The usual Stern-Gerlach experiment measures the  $\mathbf{n}$  component of the neutron spin by sending it through an inhomogeneous magnetic field  $\mathbf{B}_1(x)$  along the  $\mathbf{n}$  direction. If the neutron is polarised parallel to  $\mathbf{n}$ , it is deflected in one direction and if it is antiparallel, it suffers the opposite deflection. A general spin state of the neutron is a superposition of these two states. A neutron polarised in such a state will be deflected one way or another with probabilities determined by its overlap with the two basis states. A beam of neutrons polarised in a general direction will split into two beams, each of which is polarised parallel or antiparallel to  $\mathbf{n}$ . Note that if the beam was initially polarised along the  $\mathbf{n}$  direction, it will not split. The wavefunction does not collapse if it is already in an eigenstate of the quantity being measured.

The modified Stern-Gerlach experiment envisaged in ref.[1] uses an additional homogeneous, large magnetic field  $\mathbf{B}_0$ , which ‘prevents the wave

function from collapsing’. Ref.[1] claims that with this external field present, the beam of neutrons does not split and strikes the screen at one spot, whose location gives us information about the wave function. To quote “the beam clearly does not split provided the spin state is protected by a large homogeneous magnetic field in the *unknown* direction of the spin”.

Ref. [1] gives the impression that the proposed experiment enables one to measure the quantum state of a single system *without knowing what it was initially*. To quote “We do not know what  $|\Psi\rangle$  is before the measurement.” The purpose of this comment is to point out that one cannot *learn* anything about the system since the proposed experiment cannot be performed without prior knowledge of the spin direction. How would an experimenter create a magnetic field in an *unknown* direction? Consequently the proposed experiment does not achieve what it claims to do.

We also point out that the additional homogeneous, large magnetic field  $\mathbf{B}_0$  is not necessary in order to carry out a “protective measurement”. One can do it by means of a suitably aligned standard (unmodified) Stern-Gerlach apparatus. One can ‘measure’ the wave function of a two-state spin system by physically rotating the Stern-Gerlach apparatus so that the spin wave function is in one of its eigenstates. But in order to do this we would need to know the wave function. The situation in the modified Stern-Gerlach exper-

iment proposed in reference [1] as a ‘protected’ measurement is no different. One must know the wave function *a priori* to measure it. In the language of optics, if a photon is known to be linearly polarised (let us say it got through a polaroid) it will certainly go through polaroids aligned with the initial one. This is the simplest version of a ‘protective measurement’.

If one agrees that the proposed experiment must be a set up without prior knowledge of the wave function one finds the following well known situation. Neutrons are passed through a *fixed* magnetic field  $\mathbf{B}(\mathbf{x})$ . (The decomposition of  $\mathbf{B}$  into components  $\mathbf{B}_0$  and  $\mathbf{B}_1$  made in ref.[1] is a theoretical construct which is irrelevant to the behaviour of the system. The neutron responds to the total magnetic field in which it is placed). For any fixed choice of the magnetic field  $\mathbf{B}(x)$ , there exist spin states  $|a\rangle$  and  $|b\rangle$  (the protected ones, in the language of reference [1]) of the neutron which lead to the neutron being incident with probability one at points A and B on the screen respectively. Any other spin state  $|\psi\rangle$  is a linear combination of  $|a\rangle$  and  $|b\rangle$  and from the superposition principle, will strike the screen at A or B with probabilities given by  $|\langle\psi|a\rangle|^2$  and  $|\langle\psi|b\rangle|^2$ . If A and B are distinct, a neutron beam in a general spin state *will* split. On the other hand, if A and B are the same, all states will strike the screen at the same spot. It is easy to construct magnetic field configurations for which A and B coincide. One example is a homogeneous magnetic field. A more general

example is any field [6] which has vanishing  $\nabla|\mathbf{B}|$ . For these magnetic fields, all states strike the screen at the same spot, whose location tells us nothing whatever about the initial state. This experiment would then be a protective *non* measurement.

In summary, any experiment that increases our knowledge of the state of a quantum system necessarily disturbs it. If one does not disturb the system one learns nothing more about it. These are the basic features of quantum measurement [7] which were challenged in reference [1]. The usual interpretation of the wave function is thus protected.

## References

1. Y. Aharonov, J. Anandan and L. Vaidman, *Phys. Rev. A*, **47**, 4616 (1993). Related references are [3] and [5].
2. More precisely, the wave function up to an overall phase. In the following we ignore this relatively fine distinction between a quantum mechanical state and a ray.
3. David Freedman “Theorist to the quantum mechanical wave: Get Real” *Science*, **259**, 1542 (1993).
4. Reference [1] has also been criticised for the same basic reason in [5]. However, we disagree on salient points with [5]. To minimise confusion, we restrict our remarks here to ref.[1].
5. Carlo Rovelli, High Energy Physics Bulletin board (hep-th/9304126).
6. In the example discussed in [1]  $\mathbf{B} = \mathbf{B}_0 + \mathbf{B}_1(x)$ , where  $|\mathbf{B}_0|$  is much larger than  $|\mathbf{B}_1|$ , this situation obtains if  $\mathbf{B}_0$  is orthogonal to  $\mathbf{B}_1$ .
7. P.A.M. Dirac Principles of Quantum Mechanics, (OUP, London 1958)